

PHARMACY

Baclofen	Oral	Suspension	5mg/mL
Drug Name	Route	Dosage Form	Concentration

Formula Qty: 50mL	Shelf Life: 60 days
-------------------	---------------------

Equipment needed:

Mortar/pestle, graduated cylinder, stirring rod

Auxiliary Labeling/Storage: Refrigerate (preferred) or Room Temperature; Shake Well

Directions:

- 1. Crush tablets in mortar and pestle and triturate to a fine powder.
- 2. Add a minimal amount of vehicle and mix well to form a viscous, but smooth and uniform paste.
- 3. Continue adding vehicle in geometric portions, mixing well.
- 4. Transfer to graduate.
- 5. QS to final volume with vehicle. Stir well.

Final Product Description: Thick, opaque, off-white suspension. No discernible scent. This description is for exact products and manufacturers specified below and may vary with substitutions

Ingredients		QS	Quantity	Units
Baclofen 10mg	Ivax		25	Tablet
Ora-Blend (or 1:1 Ora-Plus/Ora-Sweet)	Paddock	X	50	mL

Citations/References

Allen LV, Erickson MA. Stability of baclofen, captopril, diltiazem HCl, dipyridamole, and flecainide acetate in extemporaneously compounded oral liquids. Am J Health Sys Pharm. 1996; (53): 2073-2078.

Reviewed: 3/12/2010 TF.

Nationwide Children's Hospital Disclaimer

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.