

Pharmacy

Drug Name	Route	Dosage Form	Concentration
Vancomycin (LOW)	Ophth	Solution	31mg/mL

Formula Qty: 15mL	Shelf Life: 21 days
-------------------	---------------------

Equipment needed:

Laminar air flow hood, syringes, needles, 0.22 micron filter

Auxiliary Labels/Storage: Refrigerate; For Ophthalmic Use Only

Room Temperature Storage = 7 days

Directions:

- 1. In a laminar air flow hood, reconstitute vancomycin 500mg vial for injection with 5mL sterile water for injection to yield a 100mg/mL solution.
- 2. Remove dropper from artificial tears bottle (15mL bottle). Remove and discard 4.6mL of solution leaving 10.4mL remaining.
- 3. Withdraw 4.6mL from the reconstituted vancomycin vial into a syringe.
- 4. Attach a 0.22 micron filter to the syringe.
- 5. Transfer the vancomycin solution into the artificial tears bottle. Replace dropper, shake well to mix.

Note: The artificial tears solution tested in the primary study contained: 0.3% hydroxypropyl methylcellulose, 0.1% dextran 70, 0.01% benzalkonium chloride, and 0.05% edentate disodium.

Ingredients	QS	Quantity	Units
Vancomycin 500mg vial for Injection		4.6	mL
Sterile Water for Injection		5	mL
Artificial Tears (Natural Balance Tears)		10.4	mL

Citations/References

- 1. Fuhrman LC, Stroman RT. Stability of Vancomycin in an Extemporaneously Compounded Ophthalmic Solution. Am J Health-Syst Pharm. 1998; 55(Jul): 1386-1388.
- 2. Osborn E, Baum JL, Ernst C, et al. Stability of Ten Antibiotics in Artificial Tear Solutions. Am J Ophth. 1976; (82): 775-780.

Reviewed: 4/13/2010 JEB

Nationwide Children's Hospital Disclaimer

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.