PUBLICATIONS:

1. Kitagawa R, Mitsuki H, Okazaki T, Ogawa T. A novel DnaA protein-binding site at 94.7 min on the Escherichia coli chromosome. Mol Microbiol 19(5):1137-1147, 1996.

2. Kitagawa R, Ozaki T, Moriya S, Ogawa T. Negative control of replication initiation by a novel chromosomal locus exhibiting exceptional affinity for Escherichia coli DnaA protein. Genes Dev 12(19):3032-3043, 1998.

3 Kitagawa R, Rose AM. Components of the spindle-assembly checkpoint are essential in Caenorhabditis elegans. Nat Cell Biol 1(8):514-521, 1999.

4. Furuta T, Tuck S, Kirchner J, Koch B, Auty R, Kitagawa R, Rose AM, Greenstein D. EMB-30: an APC4 homologue required for metaphase-to-anaphase transitions during meiosis and mitosis in Caenorhabditis elegans. Mol Biol Cell 11(4):1401-1419, 2000.

5. Ozaki T, Kumaki Y, Kitagawa R, Ogawa T. Anomalous DnaA protein binding to the regulatory region of the Escherichia coli aldA gene. Microbiology 147 (Pt 1):153-159, 2001.

6. Kitagawa R*, Law E, Tang L, Rose AM. The Cdc20 homolog, FZY-1 and its interacting protein, IFY-1 are required for proper chromosome segregation in Caenorhabditis elegans. Curr Biol 12(24):2118-2123, 2002.

7. Kitagawa R, Bakkenist CJ, McKinnon PJ, Kastan MB. Phosphorylation of SMC1 is the critical downstream event in the ATM-NBS1-BRCA1 pathway. Genes Dev 18(12):1423-1438, 2004.

8. Morales M, Theunissen JW, Kim CF, Kitagawa R, Kastan MB, Petrini JH. The Rad50S allele promotes ATM-dependent DNA damage responses and suppresses ATM deficiency: Implications for the Mre11 complex as a DNA damage sensor. Genes Dev 19(24):3043-3054, 2005.

9. Kitagawa R, Kastan MB. The ATM-dependent DNA damage signaling pathway. Cold Spring Harb Symp Quant Biol LXX 70:99-109, 2005.

10 Bekker-Jensen S, Lukas C, Kitagawa R, Melander F, Kastan MB, Bartek J, Lukas J. Spatial organization of the mammalian genome surveillance machinery in response to DNA strand breaks. J Cell Biol 173(2):195-206, 2006.

11. Tarailo M, Kitagawa R, Rose A. Suppressors of spindle checkpoint defect (such) mutants identify new mdf-1/MAD1 interactors in Caenorhabditis elegans. Genetics 175(4):1665-1679, 2007.

12. Watanabe S, Yamamoto TG, Kitagawa R*. Spindle assembly checkpoint gene mdf-1 regulates germ cell proliferation in response to nutrition signals in C. elegans. EMBO J27(7):1085-1096, 2008.

13. Yamamoto TG, Watanabe S, Essex A, Kitagawa R*. SPDL-1 functions as a kinetochore receptor for MDF-1 in Caenorhabditis elegans. J Cell Biol 183(2): 187-194, 2008.

14. Kitagawa R*. Key players in chromosome segregation in Caenorhabditis elegans. Front Biosci 14: 1529-1557, 2009.

15. Kitagawa R*. Spindle Assembly Checkpoint in C. elegans. Cell Cycle. 8(3): 1-7, 2009.

*: Corresponding author