Proportion of children who had completed the 4:3:1 Immunization Series¹ by their 2nd birthday

		•		
	Columbus City ^a	Columbus Suburban ^a	Ohio ^b	United States ^b
1996	60.0%	83.0%	72.6%	73.2%
2000	62.0%	83.0%	67.9%	75.3%
2004	64.0%	87.0%	80.1%	80.0%
2008	66.0%	88.0%	NA^2	NA^2

Source:

Notes:

¹ 4:3:1 series combination = 4 doses DTP or DTaP, 3 doses Polio, and 1 dose MMR

Contact for the Eight major Cities, Suburbs, and Rural Counties Kindergarten Retrospective Survey: Amy Bashforth Amy.Bashforth@ODH.Ohio.gov

² For 2008, the 4:3:4 period is not tracked on Contact for Disease Control and Provention's (CDC) National Immunication Survey.

² For 2008, the 4:3:1 series is not tracked on Centers for Disease Control and Prevention's (CDC) National Immunization Survey.

2008 data are available for the 4:3:1:3 (4 doses DTP or DTaP, 3 doses Polio, 1 dose MMR, and 3 doses Hib)

Ohio = 81.2% completed by age 2

Unitd States= 75.1% completed by age 2

^a Eight Major Cities, Suburbs, and Rural Counties Kindergarten Retrospective Surveys conducted by the Ohio Department of Health Immunization Program (BIDC) and Bureau of Health Surveillance (BHSIOS), 1996, 2000, 2004, and 2008.

^b National Immunization Survey, 1996, 2000, 2004, 2008, Centers for Disease Control and Prevention http://www.cdc.gov/vaccines/stats-surv/imz-coverage.htm#nis