

CPP-ED-IP-Surgery Appendicitis Clinical Pathway Published: 10/3/2023; Last Revised: 10/3/2023



• IV pain medications for breakthrough pain or if not tolerating diet

if required inpatient Phone follow-up in 3-4 weeks



CPP-ED-IP-Surgery Appendicitis Clinical Pathway Published: 10/3/2023; Last Revised: 10/3/2023



Differential Diagnoses

- Constipation
- Gastroenteritis
- Inflammatory Bowel Disease
- Meckel's Diverticulitis
- Omental infarction
- Pyelonephritis
- Pelvic Inflammatory Disease
- Tubal Ovarian Abscess
- Ectopic Pregnancy
- Typhlitis
- Ovarian Torsion
- Ruptured ovarian cyst
- Intussusception
- Testicular Torsion
- Mesenteric Adenitis
- Kidney Stones
- Spontaneous bacterial peritonitis
- Pneumonia
- Small bowel Obstruction

Return to Acute Non-Operative Algorithm	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
---	--	--	---

Definition & Diagnosis

Topic/condition definition: Appendicitis is defined as inflammation of the appendix (small tubular organ attached to the large intestine), that causes pain in the right lower quadrant of the abdomen. Pain can also be accompanied by nausea, vomiting, anorexia and low grade fevers.

Diagnostic Criteria: Postoperative Diagnosis of Appendicitis or Confirmed Abscess on Imaging

- Preadmit Findings for admission
 - Confirmed CT or US diagnosis of appendicitis (enlarged appendix over 7 mm diameter, wall thickening over 2 mm, peri-appendiceal fat stranding, fluid-filled appendix, or free fluid).
 - Right lower quadrant abdominal pain
 - o WBC elevated
 - Confirmation by surgical team (physical exam)
- **Consider other alternate clinical problem or/or diagnosis when:** Signs/Symptoms of sepsis, Bloody stools, Oliguria, Positive Pregnancy Test, Respiratory Distress
- Differential Diagnosis

Return to Acute Non-Operative Algorithm	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
---	--	--	---

Signs & Symptoms

- Periumbilical pain that migrates to right lower quadrant
- Pain typically precedes other symptoms
- Nausea
- Vomiting
- Fever
- Diarrhea
- Low grade fever
- Rebound tenderness and/or guarding
- Rovsing's sign (right sided pain from left lower abdominal palpation)
- Obturator sign (pain with flexion and internal rotation of right hip)
- Psoas sign (pain with left side down and right hip flexion)
- Markle sign (pain with heel drop)
- Usually affects ages 10-19 years
- Children younger than 3 years have high suspicion for perforation

Types of Appendicitis

- Simple Appendicitis
 - Symptoms less than 48 hours
 - Appendix is not ruptured
 - No abscess noted
 - o No fecalith
- Complex Appendicitis
 - Usually symptoms are > 48 hours
 - o National Surgical Quality Improvement Program (NSQIP) definition
 - Hole in appendix
 - Diffuse pus
 - Fecalith (calcified stool ball outside of appendix)
 - Abscess

Return to Acute Non-Operative Algorithm

Return to AcuteReturn to ComplexOperative AlgorithmOperative Algorithm

Return to Complex Non-Operative Algorithm

Inclusion & Exclusion Criteria

Inclusion Criteria:

- CT or US confirmed diagnosis of appendicitis
- > 3 months of age

Exclusion Criteria:

- < 3 months of age</p>
- Patient with unconfirmed diagnosis of appendicitis

Admission Criteria:

• All patients with appendicitis will be admitted to Pediatric Surgery



Diagnostic Testing

- Limited Abdominal Ultrasound
- CT of abdomen and pelvis
- MRI may be considered for imaging in place of CT but appendicoliths may not be visualized.

Pre-Op Imaging

- The first choice of imaging for suspected appendicitis should be a limited abdominal ultrasound.
- Ultrasounds are reported out as one of four categories:
 - Appendix is visualized and is negative for appendicitis CONSIDERED NEGATIVE
 - Appendix not seen in its entirety but no secondary signs to indicate appendicitis CONSIDERED NEGATIVE
 - Appendix is not seen in its entirety but secondary signs indicate appendicitis CONSIDERED POSITIVE
 - Appendix is seen and positive for appendicitis CONSIDERED POSITIVE
- All radiographic findings should be correlated with the history, physical exam and blood work as appropriate.
- A CT scan should be ordered only in cases where the Ultrasound and history/physical/ blood work do not match.
- A CT scan can also be used to determine whether a drainable abscess is present; however, Radiology may drain an abscess based on Ultrasound or MRI findings alone



Appendicitis Pre-Op Protocol

Pre-Op Antibiotics

- Antibiotics should be started promptly in the ED (Including LCED) after imaging confirms the diagnosis
- Pediatric Surgery will decide if a patient meets non-operative criteria. If the patient does
 meet criteria, the patient will be given Ceftriaxone/Metronidazole. If the patient is allergic
 to Cephalosporin or has had difficulty breathing or anaphylaxis to Penicillin, give
 Ciprofloxacin/Metronidazole.
 - Ceftriaxone and Metronidazole must be given 0-24 hours prior to incision
 - Ciprofloxacin and Piperacillin-Tazobactam must be given 0-6 hours prior to incision
- If a patient goes to the OR without being given antibiotics and there is not enough time for Metronidazole to infuse, the patient will be given Piperacillin-Tazobactam in the OR to expedite surgery. If the patient is found to have complex appendicitis, the patient will be given Ceftriaxone/Metronidazole after surgery.

Return to Acute Non-Operative Algorithm	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
---	--	--	---

Treatment

Operative Treatment

- Laparoscopic Appendectomy
 - o Gold standard
- Open Appendectomy

Non-Operative Management

- Antibiotic only treatment for Simple Appendicitis
- Antibiotic treatment with delayed appendectomy
 - May be used for Complex Appendicitis with abscess and/or phlegmon
 - Appendectomy usually 6-8 weeks after treated with drainage and antibiotics
- Interventional Radiology Drainage or Aspiration of Abscess >20 cm²
 - Consider repeat imaging on POD#7 if patient continues to have abdominal pain, fevers, diarrhea, elevated WBC
 - US should be used primarily
 - CT if recommended by IR or surgical attending

IV Antibiotics

- Ceftriaxone 50 mg/kg q 24 hours (max dose 2000 mg)
- Metronidazole 30 mg/kg q 24 hours (max dose 1000 mg)
- Alternative Dosing for Medication Allergies
- Penicillin Allergy
 - Ciprofloxacin 15 mg/kg/dose q 12 hours (max 400 mg)
 - Metronidazole 30 mg/kg q 24 hours (max dose 1000 mg)
- Metronidazole Allergy
 - Piperacillin-Tazobactam 100 mg/kg/dose q 8 hours (max dose 4000mg)

Oral Antibiotics

- Amoxicillin/ Clavulanate 22.5 mg/kg/dose q 12 hours (max dose 875 mg)
- Alternative Dosing
 - Ciprofloxacin 15 mg/kg/dose q12 hours (max dose 500 mg)
 - Metronidazole 10 mg/kg/dose q 8 hours (max dose 500 mg)

Return to Acute

Operative Algorithm

TPN Recommendations

Return to Acute

Non-Operative

<u>Alaorithm</u>

• May order PICC /TPN if pt NPO for > 10 days, not including time pt was NPO preoperatively

Return to Complex

Operative Algorithm

Return to Complex

Non-Operative

<u>Algorithm</u>

CPP-ED-IP-Surgery Appendicitis Clinical Pathway Published: 10/3/2023; Last Revised: 10/3/2023

Complications

- Sepsis
- Peritonitis
- Small Bowel Obstruction
- Pleural effusions
- Surgical Site Infection
- Intraabdominal abscess
- Enterocutaneous Fistula
- Clostridium Difficile infection

Return to Acute Non-Operative Algorithm	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
---	--	--	---

Monitoring, Signs of Deterioration & Escalation of Care

Monitoring protocol/process:

- Vitals signs q 4 hours (Temp, HR, RR, BP, and pOx)
- Strict I&Os
- Monitor return of bowel function (+flatus, +stool)
- Monitor diarrhea
- Monitor abdominal distention

Severity Categories/Criteria:

- Red Flags: Sepsis, Respiratory Distress, Bloody stool, Bilious emesis, Postoperative Anemia, Hypotension
- Signs and Symptoms of Sepsis: Fever, Tachycardia, Hypotension, Absent or near absent urine output, Severe pain, and Confusion

Signs of Deterioration:

- Tachycardia
- Increased abdominal pain
- Increased abdominal distention
- Peritonitis
- Anorexia
- Increased WBC (White Blood Cells)
- Increased work of breathing
- Oliguria or Anuric
- Confusion
- Fever

PICU Admission Criteria:

- Hemodynamic instability
- Inability to close abdomen postoperatively
- Respiratory distress or failure
- Sepsis

Return to Acute Non-Operative Algorithm	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
---	--	--	---

Medications

IV antibiotics

- Ceftriaxone 50 mg/kg/day IV every 24 hours (2000 mg max dose)
- Metronidazole 30 mg/kg/day IV every 24 hours (1000 mg max dose)

Alternative Antibiotics if penicillin/cephalosporin allergy

- Ciprofloxacin 10 mg/kg IV every 12 hours (400 mg max)
- Metronidazole 30 mg/kg IV every 24 hours (1000 mg max)

Alternative if Flagyl allergy

Piperacillin/Tazobactam 100 mg/kg IV every 8 hours (max 4000 mg)

Inpatient Pain Management:

- IV pain medications if not tolerating diet or pain uncontrolled with oral medications
- Ketorolac 0.5 mg/kg IV (max 30mg) every 6 hours for 12 doses initially (max 20 doses)
 - Acetaminophen IV (Ofirmev) 15 mg/kg (max 1000 mg) every 6 hours
 - Morphine 0.1 mg/kg every 3 hours as needed for pain (max 4mg, if not effective may prescribe Hydromorphone
 - o Hydromorphone (Dilaudid) 0.01-0.015 mg/kg every 2-3 hours as needed for pain
- Transition to oral medications as soon as tolerating a diet
- Acetaminophen 15 mg/kg PO (max 1000 mg) every 6 hours for 48 hours then every 6 hours as needed
- Ibuprofen 10 mg/kg PO (max 600 mg) every 6 hours for 48 hours then every 6 hours as needed
 Oxycodone 0.1 mg/kg (max 10 mg) PO Q4-6 hours as needed for severe pain

Outpatient Pain Management

- Pain Management:
 - Acetaminophen 15 mg/kg PO Q6H for 48 hours then every 4-6 hours prn
 - o Ibuprofen 10 mg/kg PO Q6H for 48 hours then every 6-8 hours prn
- Oxycodone 0.1 mg/kg PO Q6H prn ONLY if required inpatient (limit doses)

Return to Acute Non-Operative AlgorithmReturn to Acute Operative Algo	Ite Return to Complex Operative Algorithm	Return to Complex Non-Operative Algorithm
--	---	---

Discharge Criteria & Planning

Floor Discharge Criteria:

- Afebrile (temp < 100.4 F) x 24 hours
- Pain controlled with oral medications
- Tolerating regular diet: must consume 3 meals consecutively of 50% or more of typical appetite as described by caregivers. If a meal is skipped or does not take 50%, the count will restart.
- Simple appendicitis:
 - o 1 dose of IV antibiotics preoperatively and no antibiotics for home
- Complex appendicitis, operative:
 - Continue IV antibiotics until improved clinically or transition to oral prior to discharge. Treatment x 7 days total
 - CBC w/diff: if WBC ≥ 12000 then complete 7 days total, discuss with surgeon if 7 days already completed
 - CBC w/diff: if WBC < 12000 then no antibiotics for home
- Complex appendicitis with abscess drainage:
 - Continue IV antibiotics until clinically improved then transition to oral antibiotics. Treatment duration 7 days from last IR drain.
- Complex appendicitis with phlegmon or abscess not amenable to drainage:
 - Continue with IV antibiotics until clinically well and then transition to oral antibiotics. Treatment duration 10 days total

Non-Operative Return to Acute H Algorithm Operative Algorithm Operative Algorithm Operative Algorithm	Operative Algorithm	<u>Non-Operative</u> Algorithm
---	---------------------	-----------------------------------

Patient & Caregiver Education

- Helping Hands
 - o Appendectomy Simple Appendicitis HH-I-75 (Revised 2022)
 - o Appendectomy Complex Appendicitis HH-I-76 3/85 (Revised 2018)
- Footsteps to home after Ruptured Appendicitis checklist for patients and families provided by unit
- Discharge instruction template

Return to Acute Non-Operative	Return to Acute Operative Algorithm	Return to Complex Operative Algorithm	Return to Complex Non-Operative
<u>Algorithm</u>			<u>Algorithm</u>

Metrics

Pathway Goal

• Standardize criteria for post-operative management of complicated appendicitis by reducing unnecessary imaging and procedures, and duration of antibiotics.

Quality Measures

Outcome Metrics

- Reduce the number of complex appendicitis patients undergoing CT scan post-operatively.
- Reduce the number of patients undergoing percutaneous drainage of fluid collections <20 cm³ in size.
- Reduce the average duration of antibiotics for complex appendicitis post-operatively
- Reduce length of stay

Process Metrics

Clinical Pathway Utilization

Balancing Metrics

- Readmission rates within 7 days
- Delayed percutaneous drainage or prolonged length of stay for post-op complex appendicitis?



References

- 1. Baumann LM, Williams K, Oyetunji TA, Grabowski J, Lautz TB. Optimal timing of postoperative imaging for complicated appendicitis. *J Laparoendosc Adv Surg Tech A*. 2018;28(10):1248-1252. doi:10.1089/lap.2018.0121.
- 2. Chang CR, Cheng CY. Drainage of periappendiceal abscess and removal of free fecalith—extraperitoneal approach. *Int Surg.* 2014;99(4):379-383. doi:10.9738/INTSURG-D-13-00040.1.
- 3. Desai AA, Alemayehu H, Holcomb GW III, St. Peter SD. Safety of a new protocol decreasing antibiotic utilization after laparoscopic appendectomy for perforated appendicitis in children: a prospective observational study. *J Pediatr Surg.* 2015;50:912-914.
- 4. Fagenholz PJ, Peev MP, Thabet A, et al. Abscess due to perforated appendicitis: factors associated with successful percutaneous drainage. *Am J Surg.* 2016;212(4):794-798. doi:10.1016/j.amjsurg.2015.07.017.
- 5. Gordon AJ, Choi JH, Ginsburg H, Kuenzler K, Fisher J, Tomita S. Oral antibiotics and abscess formation after appendectomy for perforated appendicitis in children. *J Surg Res.* 2020;256:56-60.
- 6. Gorter RR, Meiring S, van der Lee JH, Heij HA. Intervention not always necessary in post-appendectomy abscesses in children: clinical experience in a tertiary surgical center and overview of the literature. *Eur J Pediatr.* 2016;175:1185-1191.
- 7. Hogan MJ, Marshalleck FE, Sidhu MK, et al. Quality improvement guidelines for pediatric abscess and fluid drainage. *J Vasc Interv Radiol.* 2012;23(11):1397-1402. doi:10.1016/j.jvir.2012.06.016.
- 8. Hu A, Li J, Vacek J, et al. Antibiotic resistance is common in the cultures of intraabdominal abscess drainage after appendectomy. *J Pediatr Surg.* 2022;57:102-106.
- 9. Jen J, Hwang R, Mattei P. Post-discharge antibiotics do not prevent intra-abdominal abscesses after appendectomy in children. *J Pediatr Surg.* 2023;58:258-262.
- 10. Ketha B, Stephenson KJ, Dassinger MS 3rd, Smith SD, Burford JM. Eliminating use of home oral antibiotics in pediatric complicated appendicitis. *J Surg Res.* 2021;263:151-154. doi:10.1016/j.jss.2020.12.059.
- 11. Khirallah MG, Elsossuky N, Ghazaly MO. Predictive factors of post-laparoscopic appendectomy peritoneal collection in children and adolescents with complicated appendicitis. *Afr J Paediatr Surg*. 2021;18:190-194.
- 12. Kumar RR, Kim JT, Haukoos JS, Macias LH, Dixon MR, Stamos MJ, et al. Factors affecting the successful management of intra-abdominal abscesses with antibiotics and the need for percutaneous drainage. *Dis Colon Rectum*. 2006;49(2):183-189.
- Makris GC, Malietzis G, Uberoi R. Commentary on "Society of Interventional Radiology Quality Improvement Standards for Image-Guided Percutaneous Drainage and Aspiration of Abscesses and Fluid Collections." *Cardiovasc Intervent Radiol.* 2020;43(7):1108-1109. doi:10.1007/s00270-020-02491-9.
- 14. McDaniel JD, Warren MT, Pence JC, Ey EH. Ultrasound-guided transrectal drainage of deep pelvic abscesses in children: a modified and simplified technique. *Pediatr Radiol.* 2015;45(3):435-438. doi:10.1007/s00247-014-3154-2.
- 15. Okita Y, Mohri Y, Kobayashi M, et al. Factors influencing the outcome of image-guided percutaneous drainage of intraabdominal abscess after gastrointestinal surgery. *Surgery Today*. 2013;43:1095-1102.
- 16. Okoye BO, Rampersand B, Marantos A, Abernathy LJ, Losty PD, Lloyd DA. Abscess after appendectomy in children: the role of conservative management. *Br J Surg*. 1998;85:1111-1113.
- 17. Piper HG, Derinkuyu B, Koral K, Perez EA, Murphy JT. Is it necessary to drain all postoperative fluid collections after appendectomy for perforated appendicitis? *J Pediatr Surg.* 2011;46(6):1126-1130.
- 18. Snelling CM, Poenaru D, Drover JW. Minimum postoperative antibiotic duration in advanced appendicitis in children: a review. *Pediatr Surg Int.* 2004;20(11-12):838-845. doi:10.1007/s00383-004-1280-x.
- 19. Texas Children's Hospital. Acute appendicitis/appendectomy evidence-based guideline. *Texas Children's Hospital Evidence Based Outcomes Center*. Published February 1, 2023. Accessed April 18, 2023. https://www.texaschildrens.org/sites/ default/files/uploads/documents/outcomes/standards/Appendicitis%20Guideline%20FINAL%202023.pdf.
- Van den Boom AL, de Wijkerslooth EM, van Rosmalen J, et al. Two versus five days of antibiotics after appendectomy for complex acute appendicitis (APPIC): study protocol for a randomized controlled trial. *Trials*. 2018;19(1):263. doi:10.1186/ s13063-018-2629-0.
- 21. Yousef Y, Youssef F, Homsy M, et al. Standardization of care for pediatric perforated appendicitis improves outcomes. *J Pediatr Surg.* 2017;52(12):1916-1920. doi:10.1016/j.jpedsurg.2017.08.054.



Return to Complex Operative Algorithm Return to Complex Non-Operative Algorithm

Pathway Team & Process

Pathway Development Team Leader(s): Pediatric Surgery:		Clinical Pathways Program: Medical Director – Clinical Informatics & Emergency Medicine	
			Tomara Price, MSN, CPNP – AC/PC
	Brian Kenney, MD	Dana Noffsinger, CPNP-AC	
Members:		Business & Development Manager:	
Pediatric Surgery:		Rekha Voruganti, MBOE, LSSBB	
	Dana Noffsinger, CPNP-AC	Program Coordinators:	
	Abigail Nelson, CPNP- AC	Tahje Brown, MBA	
	Sara Mansfield, MD	Tara Dinh, BS	
Interventional Radiology:		Clinical Pathway Approved	
	Mark Hogan, MD	Medical Director – Associate Chief Quality Officer, Center for Clinical Excellence:	
Pharmacy:		Ryan Bode, MD, MBOE	
	Andrew Sager, PharmD	Advisory Committee Date: September, 2023	
Infectious Disease:		Origination Date: October, 2023	
	Jeanette Tavares, MD	Next Revision Date: October, 2026	

Clinical Pathway Development

This clinical pathway was developed using the process described in the NCH Clinical Pathway Development Manual Version 6, 2022. Clinical Pathways at Nationwide Children's Hospital (NCH) are standards which provide general guidance to clinicians. Patient choice, clinician judgment, and other relevant factors in diagnosing and treating patients remain central to the selection of diagnostic tests and therapy. The ordering provider assumes all risks associates with care decisions. NCH assumes no responsibility for any adverse consequences, errors, or omissions that may arise from the use or reliance on these guidelines. NCH's clinical pathways are reviewed periodically for consistency with new evidence; however, new developments may not be represented, and NCH makes no guarantees, representations, or warranties with respect to the information provided in this clinical pathway.

Copyright © 2023. Nationwide Children's Hospital. All rights reserved. No part of this document may be reproduced, displayed, modified, or distributed in any form without the express written permission of Nationwide Children's Hospital.

For more information about our pathways and program please contact: ClinicalPathways@NationwideChildrens.org

