

**Prednisone**  
(tablet)  
**PrednisoLONE**  
(Orapred® solution)  
**PrednisoLONE**  
(Orapred ODT™ tablet)



15mg/5mL



10mg,  15mg,  30mg



10mg



20mg

Reason for taking this medicine:



- To reduce swelling in the lungs.
- To treat SEVERE coughing, wheezing, or shortness of breath.

How and when to take medicine:



- Take by mouth 1 or 2 times, as directed by your doctor.
- Take this medicine with food so it does not upset the stomach.

Possible side effects:



- Upset stomach
- Can make you hyper
- Can make you fussy
- Can make you hungry

Call the doctor or clinic if:

- Your breathing is not better after 2 days.
- You have severe stomach pain.
- You are exposed to chickenpox or measles.
- You see blood when you go to the bathroom.
- You vomit blood.
- You have questions about this medicine or how to use it.

**Important things to remember:**

- This medicine is a steroid.
- Take only as directed by your doctor.
- Talk with your doctor if you are going to have surgery while taking this medicine.
- **Use this medicine until your doctor tells you to stop**, even if you are feeling better. This is usually for 5 to 10 days.

**How to store:**

- Store in a cool dry cabinet.
- Do not store in bathroom.
- **Keep out of reach of children.**

