COMPOUND EVALUATION FORM			
Compound Name: Dipyridamole	Container-closure	Container-closure system(s): Amber Plastic Bottle	
Strength: 10 mg/mL	Preservatives: Con	Preservatives: Contained in Ora-Blend	
Dosage Form: Oral Solution	Beyond Use Date (	Beyond Use Date (compound type): 60 days	
Product Description: Opaque Suspension	Storage: Refrigera	Storage: Refrigerate (preferred) or Room Temperature	
Auxiliary Labels: Shake Well, Refrigerate, Protect From Light			
Quality control procedures (ex: pH test, etc.): None			
Ingredients:			
Dipyridamole 50 mg Tablets			
Ora-Blend			
Instructions for preparation: Calculations:			
<ol> <li>Crush tablets in a mortar and triturate to a fine powder.</li> <li>Wet powder with minimal amount of vehicle and levigate to form a viscous, but smooth and uniform paste.</li> <li>Continue to add vehicle geometrically, mixing well after each addition</li> <li>Pour into graduate</li> <li>Rinse mortar with Ora-Blend, adding rinse to graduate, until almost final volume</li> <li>QS to final volume with Ora-Blend. Stir Well.</li> </ol>			
<b>Equipment/ Supplies:</b>			
X Mortar and Pestle X S	Stirring rod	☐ Ointment slab	
X Graduated cylinder(s)	Spatula	☐ Balance	
☐ Weigh paper X	Amber plastic bottle	☐ Ointment jar	
X Counting tray □	Syringe	☐ Funnel	
References:  1. Allen LV, Erickson MA III. Stability of baclofen, captopril, diltiazem HCl, dipyridamole, and flecainide acetate in extemporaneously compounded oral liquids. Am J Health-Syst Pharm.1996; 53(Sept 15):			

Revised: Q Vo/A Swick 08/2020

2179-2184



## Nationwide Children's Hospital Disclaimer:

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

