Pharmacy Services

COMPOUND EVALUATION FORM			
Compound Name: Flecainide	Container-closure sys	Container-closure system(s): Plastic Amber Bottle	
Strength: 20 mg/mL	Preservatives: Contain	Preservatives: Contained in Ora-Blend	
Dosage Form: Oral Suspension	Beyond Use Date (compound type): 60 days		
Product Description: Opaque Suspension	Description: Opaque Suspension Storage: Refrigerate (Preferred)		
Auxiliary Labels: Shake Well; Refrigerate			
Quality control procedures (ex: pH test, etc.): None			
Ingredients: Flecainide 100 mg Tablets Ora-Blend (1:1 Ora-Plus/Ora-Sweet)			
Instructions for preparation: Calculations:			
 Crush tablets and triturate to a fine powder in mortar and pestle. Wet powder with minimal amount of vehicle and levigate to form a viscous, but smooth and uniform mass. Add remaining vehicle geometrically, mixing well after each addition. Pour into graduate. Rinse mortar with vehicle, adding rinse to graduate, until almost final volume. QS to final volume with vehicle. Stir well. 			
Equipment/ Supplies:			
X Graduated cylinder(s) ☐ Weigh paper X	X Stirring rod I Spatula X Amber (plastic/glass) bottle I Syringe	☐ Ointment slab ☐ Balance ☐ Ointment jar ☐ Funnel	
References: 1. Allen LV Jr. Erickson MA III. Stability of	of baclofen, captopril, diltiazem h	ydrochloride, dipyridamole and	
flecainide acetate in extemporaneously compounded oral liquids. Am I Health-Syst Pharm, 1996: 53			

Revised: A Swick 05/2020

(Sep15):2179-2184.



Nationwide Children's Hospital Disclaimer:

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

