Pharmacy Services

COMPOUND EVALUATION FORM			
Compound Name: Ursodiol		Container-closure system(s): Amber glass bottle	
Strength: 60 mg/mL]	Preservatives: Contained in sin	nple syrup
Dosage Form: Oral Suspension		Beyond Use Date (compound type): 35 days	
Product Description: Opaque Suspension		Storage: Refrigerated (preferred) or Room Temperature	
Auxiliary Labels: Shake well, Refrigerate			
Quality control procedures (ex: pH test, etc.): N/A			
Ingredients: Ursodiol 300 mg capsules			
Glycerin, USP Simple syrup, NF			
Instructions for preparation:		Calculations:	
1. Open capsules in glass mortar.		Ursodiol 300 mg capsules: 30 capsules	
2. Wet powder with glycerin to form a fine paste.		Glycerin: 1.5 mL	
3. Add simple syrup vehicle geometrically, mixing well after each addition.		Simple syrup: QS 150mL	
4. Pour into graduate.		Ursodiol 300 mg capsules x 30 tablets = 9,000mg	
5. Rinse mortar several times with simple syrup, adding		C 1	
to graduate.			
6. QS with simple syrup to final volume. Stir well.			
Equipment/ Supplies:			
	X Stirring rod		☐ Ointment slab
X Graduated cylinder(s)	☐ Spatula		☐ Balance
	X Amber (glas	ss) bottle	☐ Ointment jar
X Counting tray ☐ Syringe			☐ Funnel
References:			

- 1. Johnson CE, Nesbitt J. Stability of ursodiol in an extemporaneously compounded oral liquid. Am J Health Syst Pharm. 1995;52(16):1798-1800. doi:10.1093/ajhp/52.16.1798
- 2. Johnson CE, Streetman D. Stability of oral suspension of ursodiol made from tablets. Am J Health Syst Pharm. 2002;59(4):361-3. doi:10.1079/ajhp/59.4.361

Revised: A Swick 04/2021



Nationwide Children's Hospital Disclaimer:

The information provided on this site is for educational and informational purposes and may not be relied upon to direct or provide care to a patient. This material is not intended as a substitute for consulting health care professionals. The health care professionals remain solely responsible to provide and direct patient care. The information contained on this site may not be up to date due to more recent medical developments. The preparation, dispensing, and administration of pharmaceuticals is and remains the responsibility of licensed medical professional or pharmacist. Nationwide Children's Hospital assumes no liability for any injury and/or damage to persons from use of any information contained on this site.

